



PheroFIP 19

Joint Meeting of the IOBC/WPRS Working Groups

“Pheromones and other semiochemicals in integrated production” & “Integrated Protection of Fruit Crops”

20-25 January 2019

Lisbon, Portugal

Book of Abstracts





ORAL PRESENTATIONS

ABSTRACTS

SESSION 2

NEW DEVELOPMENTS IN PEST MONITORING METHODS

CHAIRPERSON: TIM BELIEN

Improved monitoring of oriental fruit moth (Lepidoptera: Tortricidae) with terpinyl acetate plus acetic acid membrane lures in Uruguayan orchards

María Valentina Mujica¹, Alan L. Knight²

¹INIA, Instituto Nacional de Investigación Agropecuaria, Estación experimental Las Brujas, Ruta 48 km 10 Canelones 90200, Uruguay. vmujica@inia.org.uy; ²USDA-ARS, Yakima Agricultural Research Laboratory, Konnowac Pass Rd. Wapato, Washington 98951, USA. alan.knight@ars.usda.gov

Abstract

The use of sex pheromones for the management of oriental fruit moth, *Grapholita molesta* (Busck), has been broadly adopted throughout its worldwide distribution in both stone (*Prunus* spp.) and pome (*Malus domestica* Borkhausen, and *Pyrus* spp.) fruits and now includes the use of hand-applied dispensers, aerosols, sprayables and attract and kill formulations. One limitation with sex pheromone-based management programs is the difficulty in monitoring pest populations with sex pheromone-baited traps. A series of studies have recently reported on the use of terpinyl acetate in a sticky trap (Cichon et al. 2013; Knight et al. 2013) and the use of a new pheromone lure that uses the combination for the three-component sex pheromone of *G. molesta* plus the sex pheromone of *Cydia pomonella* (L.) (Knight et al. 2014; Knight et al. 2016). Recent studies with other important tortricids, such as *C. pomonella* (Knight et al. 2015), and several tortricid leafrollers (Knight et al. 2017; Basoalto et al. 2017) have shown catches can be significantly increased when kairomones are combined with acetic acid. Thus, studies with *G. molesta* in Uruguay were part of a worldwide study to test several combinations of terpinyl acetate, acetic acid, and several kairomones in delta traps with sticky liners (Mujica et al. 2018). Male and female moth catches of *G. molesta* in traps baited with various septa and membrane lures were evaluated in stone and pome fruit orchards treated with sex pheromones for mating disruption from 2015 to 2017. Membrane lures were loaded with terpinyl acetate (TA), acetic acid (AA) and (Z)-3-hexenyl acetate alone or in combinations. Two septa lures were loaded with either the three-component sex pheromone blend for *G. molesta* alone or in combination with codlemone (2-PH), the sex pheromone of *C. pomonella*. A third septum lure included the combination sex pheromone blend plus pear ester, (E,Z)-2,4-ethyl decadienoate (2-PH/PE), and a fourth septum was loaded with only β -ocimene. The first study was conducted from 15 December to 18 January (2015 – 2016) in a peach orchard treated with Isomate OFM dispensers (Pacific Biocontrol) at 250/ha and was adjacent to commercial apple orchards. This study compared five lures, including OFM-PH, 2-PH, 2-PH + BO, 2-PH + TA + AA and 2-PH + TA + AA + BO. The second study was conducted during 2016-2017 and consisted of three trials. Trial 1 was conducted from 25 January to 10 February in a “Fuji” apple block treated with OFM/CM puffers (Suterra) at 2.5 units/ha. The second trial was conducted from 26 January to 16 February in peaches treated with Checkmate OFM (Suterra) at 250 dispensers per ha. The third study was conducted from 26 January to 16 February in peaches treated with the paste formulation, SPLAT OFM (ISCA Technologies) at 3.7 kg/ha. Results were consistent across studies and showed that the addition of β -ocimene or (Z)-3-hexenyl acetate did not increase moth catches. The addition of pear ester to the sex pheromone lure only marginally increased moth catches. The use of TA and AA together significantly increased moth catches compared with the use of only one of the two components. The addition of TA/AA significantly increased moth catches when combined with the 2-PH lure. The TA/AA lure also allowed traps to catch both sexes. Catch of *C. pomonella* with the 2-PH lure was comparable to the

use of codlemone. Optimization of these complex lures can likely further improve managers ability to monitor *G. molesta* and help to develop multispecies tortricid lures for use in individual traps. The complexity of combining sex pheromones with host plant, and microbial volatiles for a suite of species may be too great to create one optimized lure. Instead, the use of suboptimal lures with some sufficient attractiveness across all species might be sufficient to establish action thresholds and to reduce monitoring costs.

Key words: apple, *Cydia pomonella*, *Grapholita molesta*, mating disruption, peach

Acknowledgements

We would all like to thank Bill Lingren, Trécé Inc., Adair, OK, for providing the experimental septa and membrane lures used extensively in these trials.

References

- Basoalto, E., Hilton, R., Judd, G. J. R., Knight, A. L. El-Sayed, A. M. et al. 2017: Evaluating the use of phenylacetonitrile plus acetic acid to monitor *Pandemis pyrusana* and *Cydia pomonella* (Lepidoptera: Tortricidae) in apple. Fla. Entomol. 100: 761-766.
- Cichon, L., Fuentes-Contreras, E., Garrido, S., Lago, J., Barros-Parada, W. et al. 2013: Monitoring oriental fruit moth (Lepidoptera: Tortricidae) with sticky traps baited with terpinyl acetate and sex pheromone. J. Appl. Entomol. 137: 275-281.
- Knight, A., Basoalto, E., Hilton, R., Molinari, F, Zoller, B. et al. 2013: Monitoring oriental fruit moth (Lepidoptera: Tortricidae) with the Ajar bait trap in orchards under mating disruption. J. Appl. Entomol. 137: 650-660.
- Knight, A., Cichon, L., Lago, J, Fuentes-Contreras, E, Barros-Parada, W., et al. 2014: Monitoring oriental fruit moth and codling moth (Lepidoptera: Tortricidae) with combinations of pheromones and kairomones. J. Appl. Entomol., 138: 783-794.
- Knight, A. L., Basoalto, E., Katalin, J. & El-Sayed, A. M. 2015: A binary host plant volatile lure combined with acetic acid to monitor codling moth (Lepidoptera: Tortricidae). Environ. Entomol. 44: 1434-1440.
- Knight, A. L., Basoalto, E. & Stelinski, L. L. 2016: Variability in the efficacy of sex pheromone lures for monitoring oriental fruit moth (Lepidoptera: Tortricidae). J. Appl. Entomol. 140: 261-267.
- Knight, A. L., El-Sayed, A. M., Judd, G. J. R. & Basoalto, E. 2017: Development of 2-phenylethanol plus acetic acid lures to monitor obliquebanded leafroller (Lepidoptera: Tortricidae) under mating disruption. J. Appl. Entomol. 141: 729-739.
- Mujica, V., Preti, M., Basoalto, E., Cichon, L. Fuentes-Contreras, E. et al. 2018: Improved monitoring of oriental fruit moth (Lepidoptera: Tortricidae) with terpinyl acetate plus acetic acid membrane lures. J. Appl. Entomol. 142: 731-744.